

Product datasheet for TA504791M

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PADI4 Mouse Monoclonal Antibody [Clone ID: OTI3E1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3E1

Applications: IF, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 299-588 of human

PADI4(NP_036519) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 73.9 kDa

Gene Name: peptidyl arginine deiminase 4

Database Link: NP 036519

Entrez Gene 23569 Human

Q9UM07

Background: This gene is a member of a gene family which encodes enzymes responsible for the

conversion of arginine residues to citrulline residues. This gene may play a role in granulocyte and macrophage development leading to inflammation and immune response. [provided by


RefSeq, Jul 2008]

Synonyms: PAD; PAD4; PAD15; PD14; PD15







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PADI4 ([RC206501], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PADI4. Positive lysates [LY402202] (100ug) and [LC402202] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-PADI4 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Anti-PADI4 mouse monoclonal antibody ([TA504791]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PADI4 ([RC206501]).