

Product datasheet for TA504950

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FTCD Mouse Monoclonal Antibody [Clone ID: OTI2B12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2B12

Applications: WB

Recommended Dilution: WB 1:500~2000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human FTCD(NP_006648) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 58.7 kDa

Gene Name: formimidoyltransferase cyclodeaminase

Database Link: NP 006648

Entrez Gene 10841 Human

<u>095954</u>

Background: The protein encoded by this gene is a bifunctional enzyme that channels 1-carbon units from

formiminoglutamate, a metabolite of the histidine degradation pathway, to the folate pool. Mutations in this gene are associated with glutamate formiminotransferase deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq,

Dec 20091







Synonyms: LCHC1

Protein Pathways: Histidine metabolism, Metabolic pathways, One carbon pool by folate

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FTCD ([RC208573], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FTCD. Positive lysates [LY416500] (100ug) and [LC416500] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-FTCD monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).