

Product datasheet for TA505052M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cbl c (CBLC) Mouse Monoclonal Antibody [Clone ID: OTI3D9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3D9

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human

Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 244-474 of human

CBLC(NP 036248) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 52.3 kDa

Gene Name: Cbl proto-oncogene C

Database Link: NP 036248

Entrez Gene 23624 Human

Q9ULV8

Background: This gene encodes a member of the Cbl family of E3 ubiquitin ligases. Cbl proteins play

important roles in cell signaling through the ubiquitination and subsequent downregulation

of tyrosine kinases. Expression of this gene may be restricted to epithelial cells, and

alternatively spliced transcript variants encoding multiple isoforms have been observed for

this gene. [provided by RefSeq, Feb 2012]





Synonyms: CBL-3; CBL-SL; RNF57

Note: WB application: Only overexpression results are guaranteed, not endogenous

Protein Families: Druggable Genome

Protein Pathways: Chronic myeloid leukemia, Endocytosis, ErbB signaling pathway, Insulin signaling pathway,

Jak-STAT signaling pathway, Pathways in cancer, T cell receptor signaling pathway, Ubiquitin

mediated proteolysis

Product images:




Figure A, Western blot analysis of overexpressed lysates (25ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], NC), human CBLC plasmid ([RC205130], hCBLC), mouse CBLC plasmid ([MR207210], mCBLC) using anti-CBLC antibody [TA505052] (1:5000@1mg/ml). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:5000@1mg/ml).