

Product datasheet for **TA505100M**


EIF5A2 Mouse Monoclonal Antibody [Clone ID: OTI6G7]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI6G7
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human EIF5A2(NP_065123) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	16.6 kDa
Gene Name:	eukaryotic translation initiation factor 5A2
Database Link:	NP_065123 Entrez Gene 208691 Mouse Entrez Gene 310261 Rat Entrez Gene 56648 Human Q9GZV4
Synonyms:	EIF-5A2; eIF5AII


[View online »](#)

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY EIF5A2 (Cat# [RC206249], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-EIF5A2 (Cat# [TA505100]). Positive lysates [LY412495] (100ug) and [LC412495] (20ug) can be purchased separately from OriGene.







Figure A, Western blot analysis of overexpressed lysates (15ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], NC), human EIF5A2 plasmid ([RC206249], hEIF5A2), human EIF5A1 plasmid ([RC200770], hEIF5A1) using anti-EIF5A2 antibody [TA505100] (1:500). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000)




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-EIF5A2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Carcinoma of Human pancreas tissue using anti-EIF5A2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Anti-EIF5A2 mouse monoclonal antibody (TA505100) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY EIF5A2 (RC206249).



HEK293T cells transfected with either [RC206249] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-EIF5A2 antibody ([TA505100]), and then analyzed by flow cytometry.