

# **Product datasheet for TA505100S**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **EIF5A2 Mouse Monoclonal Antibody [Clone ID: OTI6G7]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6G7

**Applications:** FC, IF, IHC, WB

**Recommended Dilution:** WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human EIF5A2(NP\_065123) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 16.6 kDa

**Gene Name:** eukaryotic translation initiation factor 5A2

Database Link: NP 065123


Entrez Gene 208691 MouseEntrez Gene 310261 RatEntrez Gene 56648 Human

Q9GZV4

Synonyms: EIF-5A2; eIF5AII



# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY EIF5A2 (Cat# [RC206249], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-EIF5A2(Cat# [TA505100]). Positive lysates [LY412495] (100ug) and [LC412495] (20ug) can be purchased separately from OriGene.






Figure A, Western blot analysis of overexpressed lysates(15ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], NC), human EIF5A2 plasmid ([RC206249], hEIF5A2), human EIF5A1 plasmid ([RC200770], hEIF5A1) using anti-EIF5A2 antibody [TA505100](1:500). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000)




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-EIF5A2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505100])






Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-EIF5A2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505100])



Anti-EIF5A2 mouse monoclonal antibody ([TA505100]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY EIF5A2 ([RC206249]).



HEK293T cells transfected with either [RC206249] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-EIF5A2 antibody ([TA505100]), and then analyzed by flow cytometry.