

Product datasheet for TA505106BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RPC62 (POLR3C) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2H1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2H1

Applications: IF, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human POLR3C(NP_006459) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 60.4 kDa

Gene Name: RNA polymerase III subunit C

Database Link: NP 006459

Entrez Gene 74414 MouseEntrez Gene 310685 RatEntrez Gene 10623 Human

Q9BUI4

Synonyms: RPC3; RPC62

Protein Families: Transcription Factors


Protein Pathways: Cytosolic DNA-sensing pathway, Metabolic pathways, Purine metabolism, Pyrimidine

metabolism, RNA polymerase







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY POLR3C (Cat# [RC200279], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-POLR3C(Cat# [TA505106]). Positive lysates [LY401943] (100ug) and [LC401943] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from 7 different cell lines by using anti-POLR3C monoclonal antibody (1:200).




Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-POLR3C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505106])






Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-POLR3C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505106])




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-POLR3C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505106])




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-POLR3C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505106])






Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-POLR3C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505106])




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-POLR3C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505106])




Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-POLR3C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505106])






Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-POLR3C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505106])



Immunohistochemical staining of paraffinembedded Human bladder tissue within the normal limits using anti-POLR3C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505106])



Anti-POLR3C mouse monoclonal antibody ([TA505106]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY POLR3C ([RC200279]).