

Product datasheet for TA505236AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Host cell factor C2 (HCFC2) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI6C9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI6C9
Applications: IF, WB

Reactivity: WB 1:2000, IF 1:100 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human HCFC2(NP_037452) produced in HEK293T

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 86.6 kDa

Gene Name: host cell factor C2

Database Link: NP 037452

Entrez Gene 67933 MouseEntrez Gene 314704 RatEntrez Gene 29915 Human

Q9Y5Z7





Host cell factor C2 (HCFC2) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI6C9] – TA505236AM

Background: This gene encodes one of two proteins which interact with VP16, a herpes simplex virus


protein that initiates virus infection. Both the encoded protein and the original Herpes host cell factor interact with VP16 through a beta-propeller domain. The original Herpes host cell factor, however, is effective at initiating viral infection while the encoded protein is not. Transcripts of varying length due to alternative polyadenylation signals have been described.

[provided by RefSeq]. COMPLETENESS: complete on the 3' end.


Synonyms: HCF-2; HCF2

Protein Families: Transcription Factors

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HCFC2 ([RC207543], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HCFC2. Positive lysates [LY415672] (100ug) and [LC415672] (20ug) can be purchased separately from OriGene.



Anti-HCFC2 mouse monoclonal antibody ([TA505236]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY HCFC2 ([RC207543]).