

Product datasheet for TA505328

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ZNF447 (ZSCAN18) Mouse Monoclonal Antibody [Clone ID: OTI4F12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4F12
Applications: IF, IHC, WB

Recommended Dilution: WB 1:4000, IHC 1:150, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ZSCAN18(NP_076415) produced in

HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 54.6 kDa

Gene Name: zinc finger and SCAN domain containing 18

Database Link: NP 076415

Entrez Gene 65982 Human

Q8TBC5


Synonyms: ZNF447

Protein Families: Transcription Factors







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ZSCAN18 ([RC203390], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ZSCAN18. Positive lysates [LY402961] (100ug) and [LC402961] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-ZSCAN18 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505328)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-ZSCAN18 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505328)






Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-ZSCAN18 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505328)




Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-ZSCAN18 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505328)




Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-ZSCAN18 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505328)






Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-ZSCAN18 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505328)



Immunohistochemical staining of paraffinembedded Human bladder tissue within the normal limits using anti-ZSCAN18 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505328)



Anti-ZSCAN18 mouse monoclonal antibody (TA505328) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ZSCAN18 ([RC203390]).