

Product datasheet for **TA505391AM**

ETS2 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2A3]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI2A3
Applications:	IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150, IF 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 90-349 of human ETS2(NP_005230) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	52.8 kDa
Gene Name:	ETS proto-oncogene 2, transcription factor
Database Link:	NP_005230 Entrez Gene 23872 Mouse Entrez Gene 304063 Rat Entrez Gene 2114 Human P15036
Background:	ETS transcriptions factors, such as ETS2, regulate numerous genes and are involved in stem cell development, cell senescence and death, and tumorigenesis. The conserved ETS domain within these proteins is a winged helix-turn-helix DNA-binding domain that recognizes the core consensus DNA sequence GGAA/T of target genes (Dwyer et al., 2007 [PubMed 17986575]). [supplied by OMIM]




[View online »](#)

Synonyms: ETS2IT1
Protein Families: Druggable Genome, Transcription Factors
Protein Pathways: Dorso-ventral axis formation


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ETS2 ([RC207377], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ETS2. Positive lysates [LY401607] (100ug) and [LC401607] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from 2 different cell lines by using anti-ETS2 monoclonal antibody (1:200).




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])




Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])



Immunohistochemical staining of paraffin-embedded Human bladder tissue within the normal limits using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])



Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])



Anti-ETS2 mouse monoclonal antibody ([TA505391]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ETS2 ([RC207377]).