

Product datasheet for TA505391BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ETS2 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2A3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2A3

Applications: IF, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 90-349 of human

ETS2(NP_005230) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 52.8 kDa

Gene Name: ETS proto-oncogene 2, transcription factor

Database Link: NP 005230

Entrez Gene 23872 MouseEntrez Gene 304063 RatEntrez Gene 2114 Human

P15036

Background: ETS transcriptions factors, such as ETS2, regulate numerous genes and are involved in stem

cell development, cell senescence and death, and tumorigenesis. The conserved ETS domain within these proteins is a winged helix-turn-helix DNA-binding domain that recognizes the core consensus DNA sequence GGAA/T of target genes (Dwyer et al., 2007 [PubMed

17986575]). [supplied by OMIM]






Synonyms: ETS2IT1


Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Dorso-ventral axis formation

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ETS2 ([RC207377], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ETS2. Positive lysates [LY401607] (100ug) and [LC401607] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from 2 different cell lines by using anti-ETS2 monoclonal antibody (1:200).






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])




Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])






Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])




Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])






Immunohistochemical staining of paraffinembedded Human bladder tissue within the normal limits using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])



Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-ETS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA505391])



Anti-ETS2 mouse monoclonal antibody ([TA505391]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ETS2 ([RC207377]).