

## Product datasheet for **TA505394M**


### **NKIRAS1 Mouse Monoclonal Antibody [Clone ID: OTI6C11]**

#### **Product data:**


Product Type:	Primary Antibodies
Clone Name:	OTI6C11
Applications:	IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150, IF 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human NKIRAS1(NP_065078) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	21.5 kDa
Gene Name:	NFKB inhibitor interacting Ras like 1
Database Link:	<a href="#">NP_065078</a> <a href="#">Entrez Gene 69721 Mouse</a> <a href="#">Entrez Gene 305751 Rat</a> <a href="#">Entrez Gene 28512 Human</a> <a href="#">Q9NYS0</a>
Synonyms:	kappaB-Ras1; KBRAS1


[View online »](#)


## Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NKIRAS1 (Cat# [RC204248], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NKIRAS1 (Cat# [TA505394]). Positive lysates [LY412541] (100ug) and [LC412541] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-NKIRAS1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-NKIRAS1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human bladder tissue within the normal limits using anti-NKIRAS1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Anti-NKIRAS1 mouse monoclonal antibody ([TA505394]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY NKIRAS1 ([RC204248]).