

Product datasheet for TA505447

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PCDH7 Mouse Monoclonal Antibody [Clone ID: OTI2B12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2B12

Applications: WB

Recommended Dilution: WB 1:200 - 1:1000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PCDH7(NP_115833) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 132.4 kDa

Gene Name: protocadherin 7

Database Link: NP 115833

Entrez Gene 54216 MouseEntrez Gene 5099 Human

<u>O60245</u>

Background: This gene belongs to the protocadherin gene family, a subfamily of the cadherin superfamily.

The gene encodes a protein with an extracellular domain containing 7 cadherin repeats. The

gene product is an integral membrane protein that is thought to function in cell-cell

recognition and adhesion. Alternative splicing yields isoforms with unique cytoplasmic tails.

[provided by RefSeq, Jul 2008]







Synonyms: BH-Pcdh; BHPCDH; PPP1R120

Protein Families: Druggable Genome, Transmembrane

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PCDH7 ([RC223033], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PCDH7. Positive lysates [LY410093] (100ug) and [LC410093] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 1 cell line by using anti-PCDH7 monoclonal antibody at 1:200.