

Product datasheet for TA505452BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PCDH7 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2G6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2G6

Applications: IF, IHC, WB

Recommended Dilution: WB 1:200~2000, IHC 1:150, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PCDH7(NP_115833) produced in HKE293T

cell.

Formulation: PBS (pH 7.4) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 132.4 kDa

Gene Name: protocadherin 7

Database Link: NP 115833

Entrez Gene 54216 MouseEntrez Gene 5099 Human

<u>060245</u>

Background: This gene belongs to the protocadherin gene family, a subfamily of the cadherin superfamily.

The gene encodes a protein with an extracellular domain containing 7 cadherin repeats. The

gene product is an integral membrane protein that is thought to function in cell-cell

recognition and adhesion. Alternative splicing yields isoforms with unique cytoplasmic tails.

[provided by RefSeq, Jul 2008]







Synonyms: BH-Pcdh; BHPCDH; PPP1R120

Protein Families: Druggable Genome, Transmembrane

Product images:



Western blot analysis of extracts (10ug) from 8 Human tissue by using anti-PCDH7 monoclonal antibody at 1:200 (1: Testis; 2: Uterus; 3: Breast; 4: Brain; 5: Liver; 6: Ovary; 7: Thyroid gland; 8: Colon).



Western blot analysis of extracts (50ug per lane) from 2 cell lines lysates by using anti-PCDH7 monoclonal antibody([TA505452], 1:500)








Figure A, Western blot analysis of overexpressed lysates(15ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], NC), human PCDH7 plasmid ([RC213200], hPCDH7 va), human PCDH7 plasmid ([RC23033], hPCDH7 vc), human PCDH7 plasmid ([RC230683], hPCDH7 vd), mouse PCDH7 plasmid ([MR214196], mPCDH7 v1), mouse PCDH7 plasmid ([MR214198], mPCDH7 v2) using anti-PCDH7 antibody [TA505452] (1:500). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000)



Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-PCDH7 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505452])



Anti-PCDH7 mouse monoclonal antibody ([TA505452]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PCDH7 ([RC223033]).