

Product datasheet for **TA505464BM**

NDOR1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1E3]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI1E3
Applications:	IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150, IF 1:100
Reactivity:	Human, Dog, Mouse
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human NDOR1(NP_055249) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	66.6 kDa
Gene Name:	NADPH dependent diflavin oxidoreductase 1
Database Link:	NP_055249 Entrez Gene 78797 Mouse Entrez Gene 491234 Dog Entrez Gene 27158 Human Q9UHB4
Background:	This gene encodes an NADPH-dependent diflavin reductase that contains both flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD) binding domains. The encoded protein catalyzes the transfer of electrons from NADPH through FAD and FMN cofactors to potential redox partners. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2012]




[View online »](#)

Synonyms: bA350O14.9; NR1


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NDOR1 (Cat# [RC204845], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NDOR1 (Cat# [TA505464]). Positive lysates [LY415283] (100ug) and [LC415283] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-NDOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505464])




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-NDOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505464])




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-NDOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505464])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-NDOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505464])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-NDOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505464])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-NDOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505464])



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-NDOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505464])



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-NDOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505464])



Anti-NDOR1 mouse monoclonal antibody ([TA505464]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY NDOR1 ([RC204845]).