

Product datasheet for TA505532AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Tapasin Related Protein (TAPBPL) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1C9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C9
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TAPBPL(NP_060479) produced in HEK293T

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 50 kDa

Gene Name: TAP binding protein like

Database Link: NP 060479

Entrez Gene 55080 Human

Q9BX59





Tapasin Related Protein (TAPBPL) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1C9] – TA505532AM

Background: Tapasin, or TAPBP (MIM 601962), is a member of the variable-constant Ig superfamily that


links major histocompatibility complex (MHC) class I molecules to the transporter associated with antigen processing (TAP; see MIM 170260) in the endoplasmic reticulum (ER). The TAPBP gene is located near the MHC complex on chromosome 6p21.3. TAPBPL is a member of the Ig superfamily that is localized on chromosome 12p13.3, a region somewhat paralogous to the

MHC. [supplied by OMIM, Mar 2008]


Synonyms: TAPBP-R; TAPBPR

Protein Families: Druggable Genome, Transmembrane

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TAPBPL ([RC201899], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TAPBPL. Positive lysates [LY413343] (100ug) and [LC413343] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-TAPBPL monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human). Dilution: 1:2000

Tapasin Related Protein (TAPBPL) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1C9] – TA505532AM



Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-TAPBPL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505532])



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-TAPBPL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505532])