

Product datasheet for TA505574S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NME2 Mouse Monoclonal Antibody [Clone ID: OTI2F9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2F9
Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NME2(NP_001018149) produced in

HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 17.1 kDa

Gene Name: NME/NM23 nucleoside diphosphate kinase 2

Database Link: NP 001018149

Entrez Gene 18103 MouseEntrez Gene 83782 RatEntrez Gene 4831 Human

P22392

Background: Nucleoside diphosphate kinase (NDK) exists as a hexamer composed of 'A' (encoded by

NME1) and 'B' (encoded by this gene) isoforms. Multiple alternatively spliced transcript variants have been found for this gene. Read-through transcription from the neighboring upstream gene (NME1) generates naturally-occurring transcripts (NME1-NME2) that encode a fusion protein comprised of sequence sharing identity with each individual gene product.

[provided by RefSeq]






Synonyms: NDKB; NDPK-B; NDPKB; NM23-H2; NM23B; PUF


Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Metabolic pathways, Purine metabolism, Pyrimidine metabolism

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NME2 (Cat# [RC200680], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NME2(Cat# [TA505574]). Positive lysates [LY422666] (100ug) and [LC422666] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-NME2 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).