

### Product datasheet for TA505597AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **GLB1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI5H2]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5H2

Applications: FC, IF, WB

Recommended Dilution: WB 1:1000, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human GLB1(NP\_001073279) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 72.6 kDa

**Gene Name:** galactosidase beta 1

Database Link: NP 001073279

Entrez Gene 2720 Human

P16278

**Background:** This gene encodes beta-galactosidase-1, a lysosomal enzyme that hydrolyzes the terminal

beta-galactose from ganglioside substrates and other glycoconjugates. Defects in this gene are the cause of GM1-gangliosidosis and Morquio B syndrome. Multiple transcript variants

encoding different isoforms have been found for this gene. [provided by RefSeq]

**Synonyms:** EBP; ELNR1; MPS4B






**Protein Families:** Druggable Genome

**Protein Pathways:** Galactose metabolism, Glycosaminoglycan degradation, Glycosphingolipid biosynthesis -


ganglio series, Lysosome, Metabolic pathways, Other glycan degradation, Sphingolipid

metabolism

# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GLB1 ([RC200721], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GLB1.




Western blot analysis of extracts (10ug) from 6 different cell lines by using anti-GLB1 monoclonal antibody (1:200).





Anti-GLB1 mouse monoclonal antibody ([TA505597]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY GLB1 ([RC200721]).



HEK293T cells transfected with either [RC200721] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-GLB1 antibody ([TA505597]), and then analyzed by flow cytometry.