

### Product datasheet for TA505624AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Tapasin Related Protein (TAPBPL) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1D3]

#### **Product data:**

Isotype:

**Product Type:** Primary Antibodies

Clone Name: OTI1D3

Applications: WB

Reactivity: WB 1:2000
Host: WB 0:2000

Human

Mouse

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TAPBPL(NP\_060479) produced in HEK293T

cell.

IgG2b

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 50 kDa

**Gene Name:** TAP binding protein like

Database Link: NP 060479

Entrez Gene 55080 Human

Q9BX59





# Tapasin Related Protein (TAPBPL) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1D3] – TA505624AM

Background: Tapasin, or TAPBP (MIM 601962), is a member of the variable-constant Ig superfamily that


links major histocompatibility complex (MHC) class I molecules to the transporter associated with antigen processing (TAP; see MIM 170260) in the endoplasmic reticulum (ER). The TAPBP gene is located near the MHC complex on chromosome 6p21.3. TAPBPL is a member of the Ig superfamily that is localized on chromosome 12p13.3, a region somewhat paralogous to the

MHC. [supplied by OMIM, Mar 2008]

Synonyms: TAPBP-R; TAPBPR

**Protein Families:** Druggable Genome, Transmembrane

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TAPBPL ([RC201899], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TAPBPL. Positive lysates [LY413343] (100ug) and [LC413343] (20ug) can be purchased separately from OriGene.