

Product datasheet for TA505625AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GLB1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI10B2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI10B2
Applications: IF, WB

Recommended Dilution: WB 1:200~2000, IF 1:100

Reactivity: Human, Dog, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human GLB1(NP_001073279) produced in

HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 72.6 kDa

Gene Name: galactosidase beta 1

Database Link: NP 001073279

Entrez Gene 12091 MouseEntrez Gene 316033 RatEntrez Gene 403873 DogEntrez Gene

709355 MonkeyEntrez Gene 2720 Human

P16278

Background: This gene encodes beta-galactosidase-1, a lysosomal enzyme that hydrolyzes the terminal

beta-galactose from ganglioside substrates and other glycoconjugates. Defects in this gene are the cause of GM1-gangliosidosis and Morquio B syndrome. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2008]





Synonyms: EBP; ELNR1; MPS4B


Protein Families: Druggable Genome

Protein Pathways: Galactose metabolism, Glycosaminoglycan degradation, Glycosphingolipid biosynthesis -


ganglio series, Lysosome, Metabolic pathways, Other glycan degradation, Sphingolipid

metabolism


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GLB1 ([RC200721], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GLB1.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-GLB1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Immunofluorescent staining of HeLa cells using anti-GLB1 mouse monoclonal antibody ([TA505625]).