

Product datasheet for TA505871BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

THUMPD1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1H6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1H6

Applications: IF, IHC, WB

Recommended Dilution: WB 1:200~4000, IHC 1:150, IF 1:100

Reactivity: Human, Monkey, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human THUMPD1(NP_060206) produced in

HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 39.1 kDa

Gene Name: THUMP domain containing 1

Database Link: NP 060206

Entrez Gene 233802 MouseEntrez Gene 696746 MonkeyEntrez Gene 55623 Human


Q9NXG2

Synonyms: Tan1







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY THUMPD1 (Cat# [RC201773], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-THUMPD1(Cat# [TA505871]). Positive lysates [LY413587] (100ug) and [LC413587] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-THUMPD1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-THUMPD1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505871])






Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-THUMPD1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505871])




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-THUMPD1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505871])




Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-THUMPD1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505871])






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-THUMPD1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505871])



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-THUMPD1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505871])



Anti-THUMPD1 mouse monoclonal antibody ([TA505871]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY THUMPD1 ([RC201773]).





Immunofluorescent staining of HeLa cells using anti-THUMPD1 mouse monoclonal antibody ([TA505871]).