

## **Product datasheet for TA505872**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### MAPK4 Mouse Monoclonal Antibody [Clone ID: OTI2F4]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2F4

**Applications:** IF, IHC, WB

Recommended Dilution: WB 1:4000, IHC 1:150, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MAPK4(NP\_002738) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 65.7 kDa

**Gene Name:** mitogen-activated protein kinase 4

Database Link: NP 002738

Entrez Gene 54268 RatEntrez Gene 225724 MouseEntrez Gene 5596 Human

P31152

**Background:** Mitogen-activated protein kinase 4 is a member of the mitogen-activated protein kinase

family. Tyrosine kinase growth factor receptors activate mitogen-activated protein kinases which then translocate into the nucleus where it phosphorylates nuclear targets. [provided by

RefSeq, Jul 2008]

Synonyms: ERK-4; ERK4; p63-MAPK; p63MAPK; PRKM4







**Protein Families:** 

Druggable Genome, Protein Kinase


# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MAPK4 ([RC207414], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAPK4. Positive lysates [LY419131] (100ug) and [LC419131] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-MAPK4 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Anti-MAPK4 mouse monoclonal antibody (TA505872) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY MAPK4 ([RC207414]).