

Product datasheet for TA505872BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MAPK4 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2F4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2F4

Applications: IF, IHC, WB

Recommended Dilution: WB 1:4000, IHC 1:150, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MAPK4(NP_002738) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 65.7 kDa

Gene Name: mitogen-activated protein kinase 4

Database Link: NP 002738

Entrez Gene 54268 RatEntrez Gene 225724 MouseEntrez Gene 5596 Human

P31152

Background: Mitogen-activated protein kinase 4 is a member of the mitogen-activated protein kinase

family. Tyrosine kinase growth factor receptors activate mitogen-activated protein kinases which then translocate into the nucleus where it phosphorylates nuclear targets. [provided by

RefSeq, Jul 2008]


Synonyms: ERK-4; ERK4; p63-MAPK; p63MAPK; PRKM4






Protein Families: Druggable Genome, Protein Kinase


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MAPK4 ([RC207414], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAPK4. Positive lysates [LY419131] (100ug) and [LC419131] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-MAPK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505872])



Anti-MAPK4 mouse monoclonal antibody ([TA505872]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY MAPK4 ([RC207414]).