

Product datasheet for TA505881S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ICT1 (MRPL58) Mouse Monoclonal Antibody [Clone ID: OTI2F9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2F9
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ICT1(NP_001536) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 23.4 kDa

Gene Name: mitochondrial ribosomal protein L58

Database Link: NP 001536

Entrez Gene 3396 Human

Q14197

Background: The adult colon epithelium contains 3 differentiated cell types that arise from a multipotent

stem cell. Deviation from the normal maturation pathway by neoplastic transformation is thought to initiate in stem cells or their early descendants. One potential marker is ICT1 whose mRNA and protein were more highly expressed in undifferentiated than in


differentiated cells. [provided by RefSeq, Jul 2008]






Synonyms: DS-1; DS1; ICT1; MRP-L58

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ICT1 (Cat# [RC204448], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ICT1(Cat# [TA505881]). Positive lysates [LY419875] (100ug) and [LC419875] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-ICT1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505881])