

# Product datasheet for TA506174BM

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## FAK (PTK2) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4D11]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4D11
Applications: IF, IHC, WB

Recommended Dilution: WB 1:4000, IHC 1:150, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PTK2(NP\_722560) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 119.1 kDa

**Gene Name:** protein tyrosine kinase 2

Database Link: NP 722560

Entrez Gene 14083 MouseEntrez Gene 25614 RatEntrez Gene 5747 Human

005397





Background:

This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesions that form between cells growing in the presence of extracellular matrix constituents. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Activation of this gene may be an important early step in cell growth and intracellular signal transduction pathways triggered in response to certain neural peptides or to cell interactions with the extracellular matrix. Several transcript variants encoding different isoforms have been found for this gene, but the full-length natures of only three of them have been determined. [provided by RefSeq, Dec 2010]

Synonyms: FADK; FADK 1; FAK; FAK1; FRNK; p125FAK; pp125FAK; PPP1R71


**Protein Families:** Druggable Genome, Protein Kinase

Protein Pathways: Axon guidance, Chemokine signaling pathway, ErbB signaling pathway, Focal adhesion,


Leukocyte transendothelial migration, Pathways in cancer, Regulation of actin cytoskeleton,

Small cell lung cancer, VEGF signaling pathway


# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PTK2 ([RC219839], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PTK2. Positive lysates [LY403521] (100ug) and [LC403521] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell by using anti-PTK2 monoclonal antibody (1:500).




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-PTK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506174])




Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-PTK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506174])






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-PTK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506174])





Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-PTK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506174])




Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-PTK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506174])









Immunofluorescent staining of HeLa cells using anti-PTK2 mouse monoclonal antibody ([TA506174]).