

Product datasheet for **TA506179AM**

Cytochrome C Oxidase subunit VIc (COX6C) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI4B11]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI4B11
Applications:	IF, IHC, WB
Recommended Dilution:	WB 1:4000, IHC 1:150, IF 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human COX6C(NP_004365) produced in HEK293T cell.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	8.6 kDa
Gene Name:	cytochrome c oxidase subunit 6C
Database Link:	NP_004365 Entrez Gene 1345 Human P09669



[View online »](#)

Background:

Cytochrome c oxidase, the terminal enzyme of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. It is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may be involved in the regulation and assembly of the complex. This nuclear gene encodes subunit VIc, which has 77% amino acid sequence identity with mouse subunit VIc. This gene is up-regulated in prostate cancer cells. A pseudogene has been found on chromosomes 16p12. [provided by RefSeq, Jul

Synonyms:

cytochrome c oxidase subunit VIc; cytochrome c oxidase subunit VIc preprotein


Protein Families:

Transmembrane


Protein Pathways:

Alzheimer's disease, Cardiac muscle contraction, Huntington's disease, Metabolic pathways, Oxidative phosphorylation, Parkinson's disease


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY COX6C (Cat# [RC200374], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-COX6C (Cat# [TA506179]). Positive lysates [LY418028] (100ug) and [LC418028] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from a mouse cell line and 3 different mouse tissues by using anti-COX6C monoclonal antibody (1:200).




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-COX6C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506179])




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-COX6C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506179])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-COX6C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506179])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-COX6C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506179])



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-COX6C mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506179])



Anti-COX6C mouse monoclonal antibody ([TA506179]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY COX6C ([RC200374]).



Immunofluorescent staining of HeLa cells using anti-COX6C mouse monoclonal antibody ([TA506179]).