

## Product datasheet for TA506236BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### CD19 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3B10]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3B10

**Applications:** FC, IF, IHC, WB

Recommended Dilution: WB 1:4000, IHC 1:150, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD19(NP\_001761) produced in HEK293T

cell

**Formulation:** PBS (pH 7.4) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 60.9 kDa

**Gene Name:** CD19 molecule

Database Link: NP 001761

Entrez Gene 930 Human

P15391

**Background:** Lymphocytes proliferate and differentiate in response to various concentrations of different

antigens. The ability of the B cell to respond in a specific, yet sensitive manner to the various antigens is achieved with the use of low-affinity antigen receptors. This gene encodes a cell surface molecule which assembles with the antigen receptor of B lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation. [provided by RefSeq, Jul

2008]







Synonyms: B4; CVID3

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** B cell receptor signaling pathway, Hematopoietic cell lineage, Primary immunodeficiency

# **Product images:**







HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CD19 (Cat# [RC202922], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD19(Cat# [TA506236]). Positive lysates [LY400678] (100ug) and [LC400678] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts(50ug) from Raji cell lines lysates by using anti-CD19 monoclonal antibody. ([TA506236], 1:500)






Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-CD19 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 3min, [TA506236]) (1:600)




Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-CD19 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 3min, [TA506236]) (1:600)




Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-CD19 mouse monoclonal antibody. ([TA506236]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min)







Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-CD19 mouse monoclonal antibody. ([TA506236]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min)




Anti-CD19 mouse monoclonal antibody ([TA506236]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CD19 ([RC202922]).



Living HEK293T cells transfected with either CD19 ([RC202922]) overexpression plasmid or empty vector control plasmid were immunostained by either anti-CD19 antibody ([TA506236]) or a nonspecific negative control antibody ([TA180143]) and then analyzed by flow cytometry. CD19 (red) or empty vector (blue) transfected cells with anti-CD19 antibody. CD19 (green) or empty vector (aqua) transfected cells with control antibody (1:100).



Flow cytometric Analysis of living RBC-lysed human peripheral blood cells, using anti-huamn CD19 antibody ([TA506236]), (Red), compared to a nonspecific negative control antibody ([TA180144]), (green) and PBS, (blue) (1:20).



Flow cytometric Analysis of living Raji cells, using anti-CD19 antibody ([TA506236]), (Red), compared to a nonspecific negative control antibody [TA180143], (green), or PBS (blue) (1:20).