

Product datasheet for TA506294S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD33 Mouse Monoclonal Antibody [Clone ID: OTI2C1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2C1

Applications: FC, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD33(NP_001763) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 38 kDa

Gene Name: CD33 molecule

Database Link: NP 001763

Entrez Gene 945 Human

P20138

Synonyms: p67; SIGLEC-3; SIGLEC3


Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Hematopoietic cell lineage






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CD33 (Cat# [RC207023], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD33(Cat# [TA506294]). Positive lysates [LY400667] (100ug) and [LC400667] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-CD33 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506294])






Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-CD33 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506294])



Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-CD33 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506294])



Flow cytometric analysis of living 293T cells transfected with CD33 overexpression plasmid ([RC207023]), Red)/empty vector ([PS100001], Blue) using anti-CD33 antibody ([TA506294]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).