

Product datasheet for TA506319

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD33 Mouse Monoclonal Antibody [Clone ID: OTI1E10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1E10
Applications: FC, IF, WB

Recommended Dilution: WB 1:200~4000, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD33(NP_001763) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 38 kDa

Gene Name: CD33 molecule

Database Link: NP 001763

Entrez Gene 945 Human

P20138


Synonyms: p67; SIGLEC-3; SIGLEC3

Protein Families: Druggable Genome, Transmembrane


Protein Pathways: Hematopoietic cell lineage




Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD33 ([RC207023], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD33 (1:500).




Western blot analysis of extracts (35ug) from 1 cell line lysates by using anti-CD33 monoclonal antibody (1:500).



Anti-CD33 mouse monoclonal antibody (TA506319) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CD33 ([RC207023]).





Flow cytometric analysis of living 293T cells transfected with CD33 overexpression plasmid ([RC207023]), Red)/empty vector ([PS100001], Blue) using anti-CD33 antibody (TA506319). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).