

Product datasheet for TA506412M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD22 Mouse Monoclonal Antibody [Clone ID: OTI4C3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4C3
Applications: IHC, WB

Recommended Dilution: WB 1:200~2000, IHC 1:150

Reactivity: Human, Dog, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD22(NP_001762) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: CD22 molecule

Database Link: NP 001762

Entrez Gene 12483 MouseEntrez Gene 308501 RatEntrez Gene 476485 DogEntrez Gene

106994739 MonkeyEntrez Gene 933 Human

P20273


Synonyms: SIGLEC-2; SIGLEC2

Protein Families: Druggable Genome, Transmembrane


Protein Pathways: B cell receptor signaling pathway, Cell adhesion molecules (CAMs), Hematopoietic cell lineage




Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD22 ([RC216939], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD22.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-CD22 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-CD22 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.