

Product datasheet for TA506533

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

alpha Synuclein (SNCA) Mouse Monoclonal Antibody [Clone ID: OTI2F6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2F6
Applications: IHC, WB

Recommended Dilution: WB 1:4000, IHC: 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SNCA(NP_000336) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 14.3 kDa

Gene Name: synuclein alpha **Database Link:** NP 000336

Entrez Gene 20617 MouseEntrez Gene 29219 RatEntrez Gene 6622 Human

P37840

Synonyms: NACP; PARK1; PARK4; PD1


Protein Families: Druggable Genome

Protein Pathways: Alzheimer's disease, Parkinson's disease







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY SNCA (Cat# [RC210606], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SNCA (Cat# TA506533). Positive lysates [LY400127] (100ug) and [LC400127] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-SNCA mouse monoclonal antibody. (TA506533) (1:150).



Immunohistochemical staining of paraffinembedded Human adult brain tissue within the normal limits using anti-SNCA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA506533) (1:500)





Immunohistochemical staining of paraffinembedded Human embryonic cerebellum within the normal limits using anti-SNCA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA506533) (1:500)