

### **Product datasheet for TA506540AM**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Tyrosine Hydroxylase (TH) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1A12]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1A12
Applications: IF, IHC, WB

Recommended Dilution: WB 1:4000, IF 1:100, IHC: 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human TH(NP\_000351) produced in HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 55.4 kDa

**Gene Name:** tyrosine hydroxylase

Database Link: NP 000351

Entrez Gene 21823 MouseEntrez Gene 25085 RatEntrez Gene 7054 Human

P07101

**Background:** The protein encoded by this gene is involved in the conversion of tyrosine to dopamine. It is

the rate-limiting enzyme in the synthesis of catecholamines, hence plays a key role in the physiology of adrenergic neurons. Mutations in this gene have been associated with autosomal recessive Segawa syndrome. Alternatively spliced transcript variants encoding

different isoforms have been noted for this gene. [provided by RefSeq, Jul 2008]




# Tyrosine Hydroxylase (TH) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1A12] – TA506540AM

Synonyms: DYT5b; DYT14; TYH


Protein Families: Druggable Genome

**Protein Pathways:** Metabolic pathways, Parkinson's disease, Tyrosine metabolism


## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TH ([RC211218], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TH. Positive lysates [LY424777] (100ug) and [LC424777] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-TH mouse monoclonal antibody. ([TA506540])



Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-TH mouse monoclonal antibody. ([TA506540])





Anti-TH mouse monoclonal antibody ([TA506540]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY TH ([RC211218]).