

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA506542AM

Tyrosine Hydroxylase (TH) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3H3]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI3H3
Applications:	IF, WB
Recommended Dilution:	WB 1:400~4000, IF 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human TH(NP_000351) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	tyrosine hydroxylase
Database Link:	<u>NP_000351</u> <u>Entrez Gene 21823 MouseEntrez Gene 25085 RatEntrez Gene 7054 Human</u> <u>P07101</u>
Background:	The protein encoded by this gene is involved in the conversion of tyrosine to dopamine. It is the rate-limiting enzyme in the synthesis of catecholamines, hence plays a key role in the physiology of adrenergic neurons. Mutations in this gene have been associated with autosomal recessive Segawa syndrome. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq, Jul 2008]
Synonyms:	DYT5b; DYT14; TYH



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Tyrosine Hydroxylase (TH) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3H3] – TA506542AM

Protein Families:

170 -130 -100 -

70

55 -

40 -

35 -


15

Druggable Genome

Protein Pathways:

Metabolic pathways, Parkinson's disease, Tyrosine metabolism


Product images:



HepG2 HeLa SVT2 A549 COS7 Jurkat MDCK PC12 MCF7

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TH ([RC211218], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TH ([TA506542]). Positive lysates [LY424777] (100ug) and [LC424777] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-TH monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).




Anti-TH mouse monoclonal antibody ([TA506542]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY TH ([RC211218]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US







Immunofluorescent staining of HeLa cells using anti-TH mouse monoclonal antibody ([TA506542]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US