

Product datasheet for **TA506751S**


ARG2 Mouse Monoclonal Antibody [Clone ID: OTI3G5]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI3G5
Applications:	IF, IHC, WB
Recommended Dilution:	WB 1:400~4000, IHC 1:150, IF 1:100
Reactivity:	Human, Dog, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human ARG2(NP_001163) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	36 kDa
Gene Name:	arginase 2
Database Link:	NP_001163 Entrez Gene 29215 Rat Entrez Gene 480364 Dog Entrez Gene 384 Human P78540
Synonyms:	arginase; arginase 2; kidney arginase; L-arginine amidinohydrolase; L-arginine ureahydrolase; nonhepatic arginase; type II
Protein Pathways:	Arginine and proline metabolism, Metabolic pathways




[View online »](#)

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ARG2 (Cat# [RC206756], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ARG2(Cat# [TA506751]). Positive lysates [LY420091] (100ug) and [LC420091] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ARG2 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).




Western blot analysis of extracts (10ug) from 3 different mouse tissues by using anti-ARG2 monoclonal antibody (1:200).




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-ARG2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506751])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-ARG2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506751])



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-ARG2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506751])



Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-ARG2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506751])



Anti-ARG2 mouse monoclonal antibody ([TA506751]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ARG2 ([RC206756]).