

Product datasheet for TA506768

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

BID Mouse Monoclonal Antibody [Clone ID: OTI3C8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3C8

Applications: IF, IHC, WB

Recommended Dilution: WB 1:400~4000, IHC 1:150, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human BID(NP_001187) produced in HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: BH3 interacting domain death agonist

Database Link: NP 001187

Entrez Gene 637 Human

P55957

Background: This gene encodes a death agonist that heterodimerizes with either agonist BAX or

antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH-terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but the full-length nature of some variants has not been defined. [provided by

RefSeq, Jul 2008]

Synonyms: FP497






Protein Families: Druggable Genome


Protein Pathways: Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Apoptosis, Natural killer cell

mediated cytotoxicity, p53 signaling pathway, Pathways in cancer, Viral myocarditis

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BID ([RC207261], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BID. Positive lysates [LY420074] (100ug) and [LC420074] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-BID monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).





Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-BID mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Anti-BID mouse monoclonal antibody (TA506768) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY BID ([RC207261]).