

Product datasheet for TA506867AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UPRT Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2D10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2D10
Applications: IF, IHC, WB

Recommended Dilution: WB 1:4000, IHC 1:150, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human UPRT(NP_659489) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 33.6 kDa

Gene Name: uracil phosphoribosyltransferase homolog

Database Link: NP 659489

Entrez Gene 139596 Human

Q96BW1

Synonyms: FUR1; UPP


Protein Families: Druggable Genome

Protein Pathways: Metabolic pathways, Pyrimidine metabolism







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY UPRT ([RC205030], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UPRT. Positive lysates [LY408068] (100ug) and [LC408068] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-UPRT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506867])




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-UPRT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506867])






Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-UPRT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506867])



Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-UPRT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506867])



Anti-UPRT mouse monoclonal antibody ([TA506867]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY UPRT ([RC205030]).