

## Product datasheet for **TA506870BM**

### ICAM1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2H4]

#### Product data:


|                         |   |
|-------------------------|---|
| Product Type:           | Primary Antibodies  |
| Clone Name:             | OTI2H4  |
| Applications:           | FC, IF, IHC, WB   |
| Recommended Dilution:   | WB 1:2000, IHC 1:150  |
| Reactivity:             | Human   |
| Host:                   | Mouse   |
| Isotype:                | IgG1  |
| Clonality:              | Monoclonal  |
| Immunogen:              | Full length human recombinant protein of human ICAM1(NP_000192) produced in HEK293T cell.   |
| Formulation:            | PBS (pH 7.3) containing 1% BSA, 50% glycerol.   |
| Concentration:          | 0.5 mg/ml   |
| Purification:           | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)   |
| Conjugation:            | HRP   |
| Storage:                | Store at -20°C as received.   |
| Stability:              | Stable for 12 months from date of receipt.  |
| Predicted Protein Size: | 55.2 kDa  |
| Gene Name:              | intercellular adhesion molecule 1   |
| Database Link:          | <a href="#">NP_000192</a><br><a href="#">Entrez Gene 3383 Human</a><br><a href="#">P05362</a>   |
| Background:             | This gene encodes a cell surface glycoprotein which is typically expressed on endothelial cells and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also exploited by Rhinovirus as a receptor. [provided by RefSeq, Jul 2008] |
| Synonyms:               | BB2; CD54; P3.58  |
| Protein Families:       | Druggable Genome, ES Cell Differentiation/IPS, Transmembrane  |




[View online »](#)

**Protein Pathways:** Cell adhesion molecules (CAMs), Leukocyte transendothelial migration, Natural killer cell mediated cytotoxicity, Viral myocarditis


**Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ICAM1 (Cat# [RC200714], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ICAM1 (Cat# [TA506870])(1:500).




Western blot analysis of extracts (35ug) from 2 different cell lines and human spleen tissue by using anti-ICAM1 monoclonal antibody (1:100).




Western blot analysis of extracts (35ug) from A549 cells (-), HUVEC cells (-) and A549 cells treated with 20ng/ml TNF- $\alpha$  for 24h (+), using anti-ICAM1 monoclonal antibody (1:100).




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-ICAM1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506870]) (1:150)




Immunofluorescent staining of living A549 cells treated with 20ng/ml TNF- $\alpha$  for 24h using anti-ICAM1 mouse monoclonal antibody ([TA506870], green, upper left; merged, upper right) or untreated A549 cells ( merged, lower right). Cell nuclei were stained with DAPI (blue, lower left) (1:100).



Immunofluorescent staining of living HUVEC cells treated with 20ng/ml TNF-a for 5h using anti-ICAM1 mouse monoclonal antibody ([TA506870], green, upper left; merged, upper right) or untreated HUVEC cells ( merged, lower right). Cell nuclei were stained with DAPI (blue, lower left) (1:100).



HEK293T cells transfected with either [RC200714] overexpress plasmid (Red), compared to an IgG isotype control, (Green) or empty vector control plasmid (Blue) were immunostained by anti-ICAM1 antibody ([TA506870]), and then analyzed by flow cytometry (1:100).



Flow cytometric Analysis of living HUVEC cells, using anti-ICAM1 antibody ([TA506870]), (Red), compared to an IgG isotype control, (green), and negative control (PBS), (Blue) (1:100).