

Product datasheet for TA506933

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL10RA Mouse Monoclonal Antibody [Clone ID: OTI1C12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C12

Applications: WB

Recommended Dilution: WB 1:1000

Reactivity: Human Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IL10RA(NP_001549) produced in HEK293T

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 60.8 kDa

Gene Name: interleukin 10 receptor subunit alpha

Database Link: NP 001549

Entrez Gene 3587 Human

Q13651

IL10RA Mouse Monoclonal Antibody [Clone ID: OTI1C12] - TA506933

Background: The protein encoded by this gene is a receptor for interleukin 10. This protein is structurally

related to interferon receptors. It has been shown to mediate the immunosuppressive signal of interleukin 10, and thus inhibits the synthesis of proinflammatory cytokines. This receptor is reported to promote survival of progenitor myeloid cells through the insulin receptor

substrate-2/PI 3-kinase/AKT pathway. Activation of this receptor leads to tyrosine


phosphorylation of JAK1 and TYK2 kinases. Two transcript variants, one protein-coding and the other not protein-coding, have been found for this gene. [provided by RefSeq, Jan 2009]

Synonyms: CD210; CD210a; CDW210A; HIL-10R; IL-10R1; IL10R

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IL10RA ([RC207020], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IL10RA. Positive lysates [LY400596] (100ug) and [LC400596] (20ug) can be purchased separately from OriGene.