

## Product datasheet for TA507108AM

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## COX4NB (EMC8) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2A4]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI2A4
Applications: IHC, WB

**Reactivity:** WB 1:4000, IHC 1:150 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human COX4NB(NP\_006058) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 23.6 kDa

**Gene Name:** ER membrane protein complex subunit 8

Database Link: NP 006058

Entrez Gene 18117 MouseEntrez Gene 361425 RatEntrez Gene 10328 Human


043402

Synonyms: C16orf2; C16orf4; COX4NB; FAM158B; NOC4






## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY COX4NB ([RC200296], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-COX4NB. Positive lysates [LY416880] (100ug) and [LC416880] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-COX4NB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA507108])






Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-COX4NB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA507108])




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-COX4NB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA507108])




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-COX4NB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA507108])






Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-COX4NB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA507108])




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-COX4NB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA507108])




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-COX4NB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA507108])







Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-COX4NB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA507108])




Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-COX4NB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA507108])



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-COX4NB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA507108])







Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-COX4NB mouse monoclonal antibody. (Heatinduced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA507108])