

Product datasheet for TA507156M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Fibrinogen gamma chain (FGG) Mouse Monoclonal Antibody [Clone ID: OTI4B8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4B8

Applications: WB

Recommended Dilution: WB 1:4000

Reactivity: Human Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human FGG(NP_000500) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 46.4 kDa

Gene Name: fibrinogen gamma chain

Database Link: NP 000500

Entrez Gene 2266 Human

P02679



Fibrinogen gamma chain (FGG) Mouse Monoclonal Antibody [Clone ID: OTI4B8] - TA507156M

Background:


The protein encoded by this gene is the gamma component of fibrinogen, a blood-borne glycoprotein comprised of three pairs of nonidentical polypeptide chains. Following vascular injury, fibrinogen is cleaved by thrombin to form fibrin which is the most abundant component of blood clots. In addition, various cleavage products of fibrinogen and fibrin regulate cell adhesion and spreading, display vasoconstrictor and chemotactic activities, and are mitogens for several cell types. Mutations in this gene lead to several disorders, including dysfibrinogenemia, hypofibrinogenemia and thrombophilia. Alternative splicing results in two transcript variants encoding different isoforms. [provided by RefSeq, Jul

Synonyms: fibrinogen; fibrinogen gamma chain; gamma chain; gamma polypeptide

Protein Families: Druggable Genome, Secreted Protein, Transmembrane

Protein Pathways: Complement and coagulation cascades

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FGG ([RC202770], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FGG. Positive lysates [LY400173] (100ug) and [LC400173] (20ug) can be purchased separately from OriGene.