

Product datasheet for TA507215

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

nttps://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SPHK1 Mouse Monoclonal Antibody [Clone ID: OTI1A6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1A6

Applications: IF, IHC, WB

Recommended Dilution: WB 1:400~4000, IHC 1:150, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SPHK1(NP_068807) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 43.8 kDa

Gene Name: sphingosine kinase 1

Database Link: NP 068807

Entrez Gene 8877 Human

Q9NYA1





Background:

The protein encoded by this gene catalyzes the phosphorylation of sphingosine to form sphingosine-1-phosphate (S1P), a lipid mediator with both intra- and extracellular functions. Intracellularly, S1P regulates proliferation and survival, and extracellularly, it is a ligand for cell surface G protein-coupled receptors. This protein, and its product S1P, play a key role in TNF-alpha signaling and the NF-kappa-B activation pathway important in inflammatory, antiapoptotic, and immune processes. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2011]


Synonyms: SPHK


Protein Families: Druggable Genome

Protein Pathways: Calcium signaling pathway, Fc gamma R-mediated phagocytosis, Metabolic pathways,

Sphingolipid metabolism, VEGF signaling pathway

Product images:






HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY SPHK1 (Cat# [RC203398], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SPHK1(Cat# TA507215). Positive lysates [LY411847] (100ug) and [LC411847] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (30ug per lane) from 2 cell lines lysates by using anti-SPHK1 monoclonal antibody (TA507215, 1:2000).






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-SPHK1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-SPHK1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Anti-SPHK1 mouse monoclonal antibody (TA507215) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY SPHK1 ([RC203398]).





Immunofluorescent staining of HeLa cells using anti-SPHK1 mouse monoclonal antibody (TA507215).