

Product datasheet for TA507262S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TXNDC5 Mouse Monoclonal Antibody [Clone ID: OTI1C3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C3
Applications: IHC, WB

Recommended Dilution: WB 1:400~4000, IHC 1:150

Reactivity: Human, Monkey, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TXNDC5(NP_071368) produced in HEK293T

cel

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 43.5 kDa

Gene Name: thioredoxin domain containing 5

Database Link: NP 071368

Entrez Gene 105245 MouseEntrez Gene 695213 MonkeyEntrez Gene 81567 Human

Q8NBS9

Background: This gene encodes a protein-disulfide isomerase. Its expression is induced by hypoxia and its

role may be to protect hypoxic cells from apoptosis. Alternative splicing results in multiple transcript variants. Read-through transcription also exists between this gene and the neighboring upstream MUTED (muted homolog) gene. [provided by RefSeq, Dec 2010]


Synonyms: ERP46, Hcc-2, UNQ364, EndoPDI, MGC3178; thioredoxin domain containing 5






Protein Families: Druggable Genome

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TXNDC5 ([RC208568], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TXNDC5.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-TXNDC5 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-TXNDC5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA507262])



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-TXNDC5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA507262])



Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-TXNDC5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA507262])