

Product datasheet for **TA590602**

TIE1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	WB 1:5000~20000, ELISA 1:100-1:2000
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	DNA immunization. This antibody is specific for the Middle Region of the target protein.
Formulation:	20 mM Potassium Phosphate, 150 mM Sodium Chloride, pH 7.0
Concentration:	0.98mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	tyrosine kinase with immunoglobulin like and EGF like domains 1
Database Link:	NP_005415 Entrez Gene 7075 Human P35590



[View online »](#)

Background:

TIE1/TIE (tyrosine kinase with Ig and EGF homology domains 1) and TIE2/Tek define a new class of the receptor tyrosine kinase (RTK) subfamily with unique structural characteristics: two immunoglobulin like domains flanking three epidermal growth factor (EGF) like domains followed by three fibronectin type III like repeats in the extracellular region and a split tyrosine kinase domain in the cytoplasmic region. Human TIE1 cDNA encodes a 1138 amino acid residue precursor protein with a putative signal peptide, an extracellular domain, and a cytoplasmic domain. Human TIE1/Fc, a disulfide linked homodimeric protein, has a calculated molecular mass of approximately 107 kDa. Due to glycosylation, the protein migrates to approximately 160 kDa in SDS PAGE under reducing conditions. TIE1 and TIE2, expressed primarily on endothelial and hematopoietic progenitor cells, play important roles in angiogenesis, vasculogenesis, and hematopoiesis. In developing vascular endothelial cells, TIE1 and TIE2 are specifically expressed. Two ligands that bind TIE have been identified, angiopoietin 1 and angiopoietin 2. Based on gene targeting studies, the in vivo functions of TIE1 are related to endothelial cell differentiation. The receptor tyrosine kinase TIE also plays a role in the survival and integrity of the endothelium.

Synonyms:


JTK14; LMPHM11; TIE

Note:


This antibody was generated by SDIX's Genomic Antibody Technology® (GAT). [Learn about GAT](#)

Protein Families:

Druggable Genome, Protein Kinase, Transmembrane

Product images:

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TIE1 (Cat# [RC207771], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TIE1 (Cat# TA590602). Positive lysates [LY417318] (100ug) and [LC417318] (20ug) can be purchased separately from OriGene.



ELISA: TIE1 Antibody