

## **Product datasheet for TA591053**

## Troduct datasfreet for TASS 1033

SP1 Rabbit Monoclonal Antibody [Clone ID: OTIR5H7]

## Product data:

**Product Type:** Primary Antibodies

Clone Name: OTIR5H7
Applications: IHC, WB

Recommended Dilution: WB1:500, IHC 1:100 to 1:200

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Monoclonal

**Immunogen:** Full length protein

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1mg/ml

**Purification:** Purified from cell culture supernatant by affinity chromatography (protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

Predicted Protein Size: 86 kDa

**Gene Name:** Sp1 transcription factor

Database Link: Entrez Gene 6667 Human

P08047

**Background:** The protein encoded by this gene is a zinc finger transcription factor that binds to GC-rich

motifs of many promoters. The encoded protein is involved in many cellular processes; including cell differentiation; cell growth; apoptosis; immune responses; response to DNA

damage; and chromatin remodeling. Post-translational modifications such as

phosphorylation; acetylation; glycosylation; and proteolytic processing significantly affect the activity of this protein; which can be an activator or a repressor. Three transcript variants

encoding different isoforms have been found for this gene.




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (lane 1) or pCMV6-ENTRY SP1 ([RC209154], lane 2) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SP1 (TA591053, 1:500).



Immunohistochemical staining of paraffinembedded normal breast using anti-SP1 C/N TA591053 clone OTI5H7 rabbit monclonal antibody. HIER Accel buffer pH8.7 ([B22-125]) at 110C for 10 min, anti-SP1 diluted to 1:100. Detection was done with Polink1 Rabbit C/N [D13-18] with DAB Kit. Image 40x magnification.