

Product datasheet for TA591066

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Thyroxine Binding Globulin (SERPINA7) Rabbit Monoclonal Antibody [Clone ID: OTIR10C9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTIR10C9

Applications: WB

Recommended Dilution: WB1:500
Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Monoclonal

Immunogen: Full length protein

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1mg/ml

Purification: Purified from cell culture supernatant by affinity chromatography (protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

Stability: Stable for 12 months from date of receipt

Predicted Protein Size: 46 kDa

Gene Name: serpin family A member 7

Database Link: Entrez Gene 6906 Human

P05543




Thyroxine Binding Globulin (SERPINA7) Rabbit Monoclonal Antibody [Clone ID: OTIR10C9] – TA591066

Background:

There are three proteins including thyroxine-binding globulin (TBG), transthyretin and albumin responsible for carrying the thyroid hormones thyroxine (T4) and 3,5,3'-triiodothyronine (T3) in the bloodstream. This gene encodes the major thyroid hormone transport protein, TBG, in serum. It belongs to the serpin family in genomics, but the protein has no inhibitory function like many other members of the serpin family. Mutations in this gene result in TGB deficiency, which has been classified as partial deficiency, complete deficiency, and excess, based on the level of serum TBG. Alternatively spliced transcript variants encoding different isoforms have been found, but the full-length nature of these variants has not been determined.

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (lane 1, 5ug) or SerpinA7 protein (lane 2, 2ug). Samples were separated by SDS-PAGE and immunoblotted with anti-SerpinA7 (TA591066, 1:500).