

## Product datasheet for TA592032S

## AMH Rabbit Monoclonal Antibody [Clone ID: OTIR5C9]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTIR5C9
Applications:	ELISA
Recommended Dilution:	ELISA 1:5000-10000
Reactivity:	Human
Host:	Rabbit
lsotype:	IgG
Clonality:	Monoclonal
Immunogen:	Recombinant protein (26-560aa) of human AMH produced in HEK297
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Predicted Protein Size:	59.17 kDa
Gene Name:	anti-Mullerian hormone
Database Link:	<u>NP_000470</u> <u>Entrez Gene 268 Human</u> <u>P03971</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	AMH Rabbit Monoclonal Antibody [Clone ID: OTIR5C9] – TA592032S
Background:	This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate N- and C-terminal cleavage products that homodimerize and associate to form a biologically active noncovalent complex. This complex binds to the anti-Mullerian hormone receptor type 2 and causes the regression of Mullerian ducts in the male embryo that would otherwise differentiate into the uterus and fallopian tubes. This protein also plays a role in Leydig cell differentiation and function and follicular development in adult females. Mutations in this gene result in persistent Mullerian duct syndrome. [provided by RefSeq, Jul 2016]
Synonyms:	MIF; MIS
Protein Familie	s: Druggable Genome, Secreted Protein
Protein Pathwa	<b>ys:</b> Cytokine-cytokine receptor interaction, TGF-beta signaling pathway

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US