

## Product datasheet for **TA592039S**

### AMH Rabbit Monoclonal Antibody [Clone ID: OTIR4G8]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTIR4G8
Applications:	ELISA
Recommended Dilution:	ELISA 1:5000-10000
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Monoclonal
Immunogen:	Recombinant protein (26-560aa) of human AMH produced in HEK296
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Predicted Protein Size:	59.17 kDa
Gene Name:	anti-Mullerian hormone
Database Link:	<a href="#">NP_000470</a> <a href="#">Entrez Gene 268 Human P03971</a>



[View online »](#)

<b>Background:</b>	This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate N- and C-terminal cleavage products that homodimerize and associate to form a biologically active noncovalent complex. This complex binds to the anti-Mullerian hormone receptor type 2 and causes the regression of Mullerian ducts in the male embryo that would otherwise differentiate into the uterus and fallopian tubes. This protein also plays a role in Leydig cell differentiation and function and follicular development in adult females. Mutations in this gene result in persistent Mullerian duct syndrome. [provided by RefSeq, Jul 2016]
<b>Synonyms:</b>	MIF; MIS
<b>Protein Families:</b>	Druggable Genome, Secreted Protein
<b>Protein Pathways:</b>	Cytokine-cytokine receptor interaction, TGF-beta signaling pathway