

Product datasheet for **TA592781S**

glucose 6 phosphatase, catalytic subunit (G6PC) Rabbit Monoclonal Antibody [Clone ID: OTIR2H10]

Product data:

| | |
|-------------------------|--|
| Product Type: | Primary Antibodies |
| Clone Name: | OTIR2H10 |
| Applications: | IHC, WB |
| Recommended Dilution: | WB 1:2000, IHC 1:300-1:1500 |
| Reactivity: | Human |
| Host: | Rabbit |
| Isotype: | IgG |
| Clonality: | Monoclonal |
| Immunogen: | Synthetic peptide (the amino acid sequence is considered to be commercially sensitive) within Human G6PC (NP_000142). The exact sequence is proprietary. |
| Formulation: | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. |
| Concentration: | 1 mg/ml |
| Purification: | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |
| Conjugation: | Unconjugated |
| Storage: | Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws. |
| Predicted Protein Size: | 40.5 kDa |
| Gene Name: | glucose-6-phosphatase catalytic subunit 1 |
| Database Link: | NP_000142 Entrez Gene 2538 Human P35575 |



[View online »](#)

Background:

Glucose-6-phosphatase (G6Pase) is a multi-subunit integral membrane protein of the endoplasmic reticulum that is composed of a catalytic subunit and transporters for G6P, inorganic phosphate, and glucose. This gene (G6PC) is one of the three glucose-6-phosphatase catalytic-subunit-encoding genes in human: G6PC, G6PC2 and G6PC3. Glucose-6-phosphatase catalyzes the hydrolysis of D-glucose 6-phosphate to D-glucose and orthophosphate and is a key enzyme in glucose homeostasis, functioning in gluconeogenesis and glycogenolysis. Mutations in this gene cause glycogen storage disease type I (GSD1). This disease, also known as von Gierke disease, is a metabolic disorder characterized by severe hypoglycemia associated with the accumulation of glycogen and fat in the liver and kidneys. [provided by RefSeq, Feb 2011]

Synonyms:


G6Pase; G6PC; G6PT; GSD1; GSD1a

Protein Families:


Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

Protein Pathways:


Adipocytokine signaling pathway, Galactose metabolism, Glycolysis / Gluconeogenesis, Insulin signaling pathway, Metabolic pathways, Starch and sucrose metabolism

Product images:


IHC staining of FFPE human kidney tissue within the normal limits using anti-G6PC rabbit monoclonal antibody ([TA592781]) and Polink-2 HRP polymer detection kit ([D22-110]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 ([B04C-100]) at 120°C for 3 min. The brown stain indicates positive stain, blue is the counter stain.



IHC staining of FFPE human liver tissue within the normal limits using anti-G6PC rabbit monoclonal antibody ([TA592781]) and Polink-2 HRP polymer detection kit ([D22-110]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 ([B04C-100]) at 120°C for 3 min. The brown stain indicates positive stain, blue is the counter stain.



IHC staining of FFPE human duodenum tissue within the normal limits using anti-G6PC rabbit monoclonal antibody ([TA592781]) and Polink-2 HRP polymer detection kit ([D22-110]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 ([B04C-100]) at 120°C for 3 min. The brown stain indicates positive stain, blue is the counter stain.



Western blot analysis of overexpressed lysates from HEK293T cells transfected with empty plasmid ([PS100001], lane 1, 15ug) and human G6PC recombinant protein (lane 2, 100ng) using anti-G6PC antibody [TA592781](1:2000).