

Product datasheet for TA592980S

IL5 Rabbit Monoclonal Antibody [Clone ID: OTIR1G2]

Product data:

Product Type: Primary Antibodies Clone Name: OTIR1G2 **Applications:** WB Recommended Dilution: WB 1:2000 **Reactivity:** Human Host: Rabbit Isotype: lgG **Clonality:** Monoclonal Immunogen: Human recombinant protein fragment of Human IL5 (NP_000870) produced in Ecoli. PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. Formulation: Concentration: 1 mg/ml **Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) **Conjugation:** Unconjugated Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws. **Predicted Protein Size:** 15.2 kDa Gene Name: interleukin 5 Database Link: NP 000870 Entrez Gene 3567 Human P05113



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL5 Rabbit Monoclonal Antibody [Clone ID: OTIR1G2] – TA592980S

Background: This gene encodes a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. The encoded cytokine plays a major role in the regulation of eosinophil formation, maturation, recruitment and survival. The increased production of this cytokine may be related to pathogenesis of eosinophil-dependent inflammatory diseases. This cytokine functions by binding to its receptor, which is a heterodimer, whose beta subunit is shared with the receptors for interleukine 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene is located on chromosome 5 within a cytokine gene cluster which includes interleukin 4 (IL4), interleukin 13 (IL13), and CSF2. This gene, IL4, and IL13 may be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31. [provided by RefSeq, Jul 2013]

Synonyms:	EDF; IL-5; TRF
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	Allograft rejection, Asthma, Autoimmune thyroid disease, Cytokine-cytokine receptor interaction, Fc epsilon Rl signaling pathway, Hematopoietic cell lineage, Jak-STAT signaling pathway, T cell receptor signaling pathway

Product images:

NC hlL5 mlL5 NC hIL5 mIL5 170-130-100 -70 -55 -40 -35 — 25 -15 -10 -A:TA592980 B:TA180144

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US