

Product datasheet for TA594237

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Non Neuronal Enolase (ENO1) Rabbit Monoclonal Antibody [Clone ID: OTIR4G3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTIR4G3

Applications: WB

Recommended Dilution: WB1:2000

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment of Human ENO1 (NP_001419) produced in Ecoli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

Predicted Protein Size: 47.2KD

Gene Name: enolase 1

Database Link: NP 001419

Entrez Gene 13806 MouseEntrez Gene 2023 Human

P06733





Background:

This gene encodes alpha-enolase, one of three enolase isoenzymes found in mammals. Each isoenzyme is a homodimer composed of 2 alpha, 2 gamma, or 2 beta subunits, and functions as a glycolytic enzyme. Alpha-enolase in addition, functions as a structural lens protein (taucrystallin) in the monomeric form. Alternative splicing of this gene results in a shorter isoform that has been shown to bind to the c-myc promoter and function as a tumor suppressor. Several pseudogenes have been identified, including one on the long arm of chromosome 1. Alpha-enolase has also been identified as an autoantigen in Hashimoto encephalopathy. [provided by RefSeq, Jan 2011]

Synonyms: ENO1L1; HEL-S-17; MPB1; NNE; PPH

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Glycolysis / Gluconeogenesis, Metabolic pathways, RNA degradation

Product images:





Figure A, Western blot analysis of overexpressed lysates(15ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], NC), human ENO1v1 plasmid ([RC205494], hENO1v1), human ENO1v2 plasmid ([RC232529], hENO1v2), mouse ENO1 plasmid ([MR227149], mENO1), human ENO2 plasmid ([RC201085], hENO2), human ENO3 plasmid ([RC205684], hENO3) using anti-ENO1 antibody TA594237(1:2000). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000)



Western blot analysis of extracts (50ug per lane) from 6 cell lines lysates by using anti-ENO1 monoclonal antibody(TA594237, 1:2000)