

Product datasheet for **TA594367M**

AAV9 Rabbit Monoclonal Antibody [Clone ID: OTIR5D6]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTIR5D6
Applications:	WB
Recommended Dilution:	WB 1:2000
Reactivity:	Adeno-Associated Virus 9
Host:	Rabbit
Isotype:	IgG
Clonality:	Monoclonal
Immunogen:	AAV9 capsids
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.
Predicted Protein Size:	100KD
Background:	AAV vectors are promising delivery tools for human gene therapy. AAV is a single-stranded DNA parvovirus with a 4.7 kb genome composed of the rep and cap genes flanked by inverted terminal repeats (ITRs). The rep gene encodes non-structural proteins involved in viral replication, packaging, and genomic integration, whereas the cap gene codes for structural proteins (VP1, VP2, VP3) that assemble to form the viral capsid, which serves as the viral gene delivery vehicle.

[View online »](#)

Product images:





Figure A, Western blot analysis of overexpressed lysates(15ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], NC) and 16 different AAV capsid overexpressed plasmids using anti-AAV9 antibody [TA594367] (1:2000). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000).From the results,[TA594367] recognize AAV9 and AAVPHP.B capsid proteins.



Simple Western™ analysis of AAV9 purified from HEK293 using AAV9 Rabbit Monoclonal Antibody #[TA594367]. The virtual lane view (left) shows the target (as indicated) at 1:50 dilution of primary antibody. The corresponding electropherogram view (right) plots chemiluminescence by molecular weight along the capillary at 1:50 dilutions of primary antibody. This experiment was performed under reducing conditions on the Jess™ Simple Western instrument from ProteinSimple, a Bio-Techne brand, using the 12-230 kDa Separation Module.